


# FINAL NEET(UG)–2023 (MANIPUR EXAMINATION)

(Held On Tuesday 6<sup>th</sup> JUNE, 2023)

## PHYSICS

### Section-A (Physics)

1. The variation of susceptibility ( $\chi$ ) with absolute temperature (T) for a paramagnetic material is represented as :



2. A bullet of mass  $m$  hits a block of mass  $M$  elastically.

The transfer of energy is the maximum, when :

- (1)  $M = m$                       (2)  $M = 2m$   
 (3)  $M \ll m$                     (4)  $M \gg m$


3. The ground state energy of hydrogen atom is  $-13.6$  eV. The energy needed to ionize hydrogen atom from its second excited state will be :

- (1)  $13.6$  eV                      (2)  $6.8$  eV  
 (3)  $1.51$  eV                      (4)  $3.4$  eV

4. The escape velocity of a body on the earth surface is  $11.2$  km/s. If the same body is projected upward with velocity  $22.4$  km/s, the velocity of this body at infinite distance from the centre of the earth will be:

- (1)  $11.2\sqrt{2}$  km/s  
 (2) Zero  
 (3)  $11.2$  km/s  
 (4)  $11.2\sqrt{3}$  km/s

5. A lens is made up of 3 different transparent media as shown in figure. A point object O is placed on its axis beyond  $2f$ . How many real images will be obtained on the other side ?



- (1) 2  
 (2) 1  
 (3) No image will be formed  
 (4) 3

6. The diameter of a spherical bob, when measured with vernier callipers yielded the following values :  $3.33$  cm,  $3.32$  cm,  $3.34$  cm,  $3.33$  cm and  $3.32$  cm. The mean diameter to appropriate significant figures is :

- (1)  $3.328$  cm  
 (2)  $3.3$  cm  
 (3)  $3.33$  cm  
 (4)  $3.32$  cm

7. On the basis of electrical conductivity, which one of the following material has the smallest resistivity?

- (1) Germanium  
 (2) Silver  
 (3) Glass  
 (4) Silicon

8. The mechanical quantity, which has dimensions of reciprocal of mass ( $M^{-1}$ ) is :

- (1) angular momentum  
 (2) coefficient of thermal conductivity  
 (3) torque  
 (4) gravitational constant


9. The position of a particle is given by

$$\vec{r}(t) = 4t \vec{i} + 2t^2 \vec{j} + 5 \vec{k}$$


where  $t$  is in seconds and  $r$  in metre. Find the magnitude and direction of velocity  $v(t)$ , at  $t = 1$  s, with respect to  $x$ -axis

- (1)  $4\sqrt{2}$  ms<sup>-1</sup>,  $45^\circ$                       (2)  $4\sqrt{2}$  ms<sup>-1</sup>,  $60^\circ$   
 (3)  $3\sqrt{2}$  ms<sup>-1</sup>,  $30^\circ$                       (4)  $3\sqrt{2}$  ms<sup>-1</sup>,  $45^\circ$


10. For the given cycle, the work done during isobaric process is :



- (1) 200 J  
 (2) Zero  
 (3) 400 J  
 (4) 600 J
11. The equivalent capacitance of the arrangement shown in figure is :




- (1) 30 μF  
 (2) 15 μF  
 (3) 25 μF  
 (4) 20 μF
12. An ac source is connected in the given circuit. The value of  $\phi$  will be :



- (1) 60°  
 (2) 90°  
 (3) 30°  
 (4) 45°

13. The given circuit is equivalent to :



14. A particle moves with a velocity  $(5\hat{i} - 3\hat{j} + 6\hat{k})$  m s<sup>-1</sup> horizontally under the action of constant force  $(10\hat{i} + 10\hat{j} + 20\hat{k})$  N. The instantaneous power supplied to the particle is :

- (1) 200 W  
 (2) Zero  
 (3) 100 W  
 (4) 140 W
15. A certain wire A has resistance 81 Ω. The resistance of another wire B of same material and equal length but of diameter thrice the diameter of A will be :
- (1) 81 Ω (2) 9 Ω (3) 729 Ω (4) 243 Ω
16.  $\epsilon_0$  and  $\mu_0$  are the electric permittivity and magnetic permeability of free space respectively. If the corresponding quantities of a medium are  $2\epsilon_0$  and  $1.5\mu_0$  respectively, the refractive index of the medium will nearly be :

- (1)  $\sqrt{2}$  (2)  $\sqrt{3}$  (3) 3 (4) 2
17. The amount of elastic potential energy per unit volume (in SI unit) of a steel wire of length 100 cm to stretch it by 1 mm is (if Young's modulus of the wire =  $2.0 \times 10^{11}$  Nm<sup>-2</sup>) :
- (1)  $10^{11}$  (2)  $10^{17}$  (3)  $10^7$  (4)  $10^5$
18. The 4<sup>th</sup> overtone of a closed organ pipe is same as that of 3<sup>rd</sup> overtone of an open pipe. The ratio of the length of the closed pipe to the length of the open pipe is :
- (1) 8 : 9 (2) 9 : 7  
 (3) 9 : 8 (4) 7 : 9


19. A long straight wire of length 2 m and mass 250 g is suspended horizontally in a uniform horizontal magnetic field of 0.7 T. The amount of current flowing through the wire will be ( $g = 9.8 \text{ ms}^{-2}$ ):

- (1) 2.45 A (2) 2.25 A  
(3) 2.75 A (4) 1.75 A

20. According to Gauss law of electrostatics, electric flux through a closed surface depends on:

- (1) the area of the surface  
(2) the quantity of charges enclosed by the surface  
(3) the shape of the surface  
(4) the volume enclosed by the surface

21. A ball is projected from point A with velocity  $20 \text{ m s}^{-1}$  at an angle  $60^\circ$  to the horizontal direction. At the highest point B of the path (as shown in figure), the velocity  $v \text{ m s}^{-1}$  of the ball will be:



- (1) 20  
(2)  $10\sqrt{3}$   
(3) Zero  
(4) 10


22. Which of the following statement is not true?

- (1) Coefficient of viscosity is a scalar quantity  
(2) Surface tension is a scalar quantity  
(3) Pressure is a vector quantity  
(4) Relative density is a scalar quantity

23. A uniform electric field and a uniform magnetic field are acting along the same direction in a certain region. If an electron is projected in the region such that its velocity is pointed along the direction of fields, then the electron:

- (1) will turn towards right of direction of motion  
(2) will turn towards left of direction of motion  
(3) speed will decrease  
(4) speed will increase


24. A horizontal ray of light is incident on the right angled prism with prism angle  $6^\circ$ . If the refractive index of the material of the prism is 1.5, then the angle of emergence will be:



- (1)  $9^\circ$  (2)  $10^\circ$  (3)  $4^\circ$  (4)  $6^\circ$

25. A p-type extrinsic semiconductor is obtained when Germanium is doped with:



- (1) Antimony (2) Phosphorous  
(3) Arsenic (4) Boron



Which set of colours will come out in air for a situation shown in figure?

- (1) Yellow, Orange and Red  
(2) All  
(3) Orange, Red and Violet  
(4) Blue, Green and Yellow


27. If  $Z_1$  and  $Z_2$  are the impedances of the given circuits (a) and (b) as shown in figures, then choose the **correct** option



- (1)  $Z_1 < Z_2$  (2)  $Z_1 + Z_2 = 20 \Omega$   
(3)  $Z_1 = Z_2$  (4)  $Z_1 > Z_2$

28. The wavelength of Lyman series of hydrogen atom appears in:

- (1) visible region (2) far infrared region  
(3) ultraviolet region (4) infrared region



The above figure shows the circuit symbol of a transistor. Select the **correct** statements given below:

- (A) The transistor has two segments of p-type semiconductor separated by a segment of n-type semiconductor.  
(B) The emitter is of moderate size and heavily doped.  
(C) The central segment is thin and lightly doped.  
(D) The emitter base junction is reverse biased in common emitter amplifier circuit.


- (1) (C) and (D) (2) (A) and (D)  
(3) (A) and (B) (4) (B) and (C)

- 30.** The de Broglie wavelength associated with an electron, accelerated by a potential difference of 81 V is given by:
- (1) 13.6 nm (2) 136 nm  
(3) 1.36 nm (4) 0.136 nm
- 31.** The maximum power is dissipated for an ac in a/an:
- (1) resistive circuit (2) LC circuit  
(3) inductive circuit (4) capacitive circuit
- 32.** The maximum kinetic energy of the emitted photoelectrons in photoelectric effect is independent of :
- (1) work function of material  
(2) intensity of incident radiation  
(3) frequency of incident radiation  
(4) wavelength of incident radiation
- 33.** Two particles A and B initially at rest, move towards each other under mutual force of attraction. At an instance when the speed of A is  $v$  and speed of B is  $3v$ , the speed of centre of mass is :
- (1)  $2v$  (2) zero  
(3)  $v$  (4)  $4v$
- 34.** A charge  $Q \mu\text{C}$  is placed at the centre of a cube. The flux coming out from any one of its faces will be (in SI unit) :
- (1)  $\frac{Q}{\epsilon_0} \times 10^{-6}$  (2)  $\frac{2Q}{3\epsilon_0} \times 10^{-3}$   
(3)  $\frac{Q}{6\epsilon_0} \times 10^{-3}$  (4)  $\frac{Q}{6\epsilon_0} \times 10^{-6}$
- 35.** The viscous drag acting on a metal sphere of diameter 1 mm, falling through a fluid of viscosity 0.8 Pa s with a velocity of  $2 \text{ m s}^{-1}$  is equal to :
- (1)  $15 \times 10^{-3} \text{ N}$  (2)  $30 \times 10^{-3} \text{ N}$   
(3)  $1.5 \times 10^{-3} \text{ N}$  (4)  $20 \times 10^{-3} \text{ N}$

### Section-B (Physics)

- 36.** If  $R$  is the radius of the earth and  $g$  is the acceleration due to gravity on the earth surface. Then the mean density of the earth will be :
- (1)  $\frac{\pi R G}{12g}$  (2)  $\frac{3\pi R}{4gG}$  (3)  $\frac{3g}{4\pi R G}$  (4)  $\frac{4\pi G}{3gR}$
- 37.** A copper wire of radius 1 mm contains  $10^{22}$  free electrons per cubic metre. The drift velocity for free electrons when 10 A current flows through the wire will be (Given, charge on electron =  $1.6 \times 10^{-19} \text{ C}$ ) :
- (1)  $\frac{6.25 \times 10^4}{\pi} \text{ m s}^{-1}$  (2)  $\frac{6.25}{\pi} \times 10^3 \text{ m s}^{-1}$   
(3)  $\frac{6.25}{\pi} \text{ m s}^{-1}$  (4)  $\frac{6.25 \times 10^5}{\pi} \text{ m s}^{-1}$


- 38.** An object is mounted on a wall. Its image of equal size is to be obtained on a parallel wall with the help of a convex lens placed between these walls. The lens is kept at distance  $x$  in front of the second wall. The required focal length of the lens will be :
- (1) less than  $\frac{x}{4}$   
(2) more than  $\frac{x}{4}$  but less than  $\frac{x}{2}$   
(3)  $\frac{x}{2}$   
(4)  $\frac{x}{4}$
- 39.** If a conducting sphere of radius  $R$  is charged. Then the electric field at a distance  $r$  ( $r > R$ ) from the centre of the sphere would be, ( $V =$  potential on the surface of the sphere)
- (1)  $\frac{rV}{R^2}$  (2)  $\frac{R^2V}{r^3}$  (3)  $\frac{RV}{r^2}$  (4)  $\frac{V}{r}$
- 40.** A block of mass 2 kg is placed on inclined rough surface AC (as shown in figure) of coefficient of friction  $\mu$ . If  $g = 10 \text{ m s}^{-2}$ , the net force (in N) on the block will be :



- (1)  $10\sqrt{3}$  (2) zero  
(3) 10 (4) 20
- 41.** A container of volume  $200 \text{ cm}^3$  contains 0.2 mole of hydrogen gas and 0.3 mole of argon gas. The pressure of the system at temperature 200 K ( $R = 8.3 \text{ J K}^{-1} \text{ mol}^{-1}$ ) will be :-
- (1)  $6.15 \times 10^5 \text{ Pa}$  (2)  $6.15 \times 10^4 \text{ Pa}$   
(3)  $4.15 \times 10^5 \text{ Pa}$  (4)  $4.15 \times 10^6 \text{ Pa}$
- 42.** To produce an instantaneous displacement current of 2 mA in the space between the parallel plates of a capacitor of capacitance  $4 \mu\text{F}$ , the rate of change of applied variable potential difference  $\left(\frac{dV}{dt}\right)$  must be :
- (1) 800 V/s (2) 500 V/s  
(3) 200 V/s (4) 400 V/s
- 43.** An emf is generated by an ac generator having 100 turn coil, of loop area  $1 \text{ m}^2$ . The coil rotates at a speed of one revolution per second and placed in a uniform magnetic field of 0.05 T perpendicular to the axis of rotation of the coil. The maximum value of emf is :-
- (1) 3.14 V (2) 31.4 V (3) 62.8 V (4) 6.28 V



44. For very high frequencies, the effective impedance of the circuit (shown in the figure) will be :-



- (1)  $4\ \Omega$     (2)  $6\ \Omega$     (3)  $1\ \Omega$     (4)  $3\ \Omega$
45. A constant torque of  $100\ \text{N m}$  turns a wheel of moment of inertia  $300\ \text{kg m}^2$  about an axis passing through its centre. Starting from rest, its angular velocity after 3s is :-  
 (1)  $1\ \text{rad/s}$     (2)  $5\ \text{rad/s}$     (3)  $10\ \text{rad/s}$     (4)  $15\ \text{rad/s}$
46. The emf of a cell having internal resistance  $1\ \Omega$  is balanced against a length of 330 cm on a potentiometer wire. When an external resistance of  $2\ \Omega$  is connected across the cell, the balancing length will be :  
 (1) 220 cm    (2) 330 cm    (3) 115 cm    (4) 332 cm
47. A 1 kg object strikes a wall with velocity  $1\ \text{m s}^{-1}$  at an angle of  $60^\circ$  with the wall and reflects at the same angle. If it remains in contact with wall for 0.1 s, then the force exerted on the wall is :-  
 (1)  $30\sqrt{3}\ \text{N}$                       (2) Zero  
 (3)  $10\sqrt{3}\ \text{N}$                         (4)  $20\sqrt{3}\ \text{N}$

48. The angular momentum of an electron moving in an orbit of hydrogen atom is  $1.5\left(\frac{h}{\pi}\right)$ . The energy in the same orbit is nearly.

(1)  $-1.5\ \text{eV}$     (2)  $-1.6\ \text{eV}$     (3)  $-1.3\ \text{eV}$     (4)  $-1.4\ \text{eV}$

49. A particle is executing uniform circular motion with velocity  $\vec{v}$  and acceleration  $\vec{a}$ . Which of the following is true ?

(1)  $\vec{v}$  is a constant;  $\vec{a}$  is not a constant  
 (2)  $\vec{v}$  is not a constant;  $\vec{a}$  is not a constant  
 (3)  $\vec{v}$  is a constant;  $\vec{a}$  is a constant  
 (4)  $\vec{v}$  is not a constant;  $\vec{a}$  is a constant

50. A simple pendulum oscillating in air has a period of  $\sqrt{3}\ \text{s}$ . If it is completely immersed in non-viscous liquid, having density  $\left(\frac{1}{4}\right)^{\text{th}}$  of the material of the bob, the new period will be :-

(1)  $2\sqrt{3}\ \text{s}$                               (2)  $\frac{2}{\sqrt{3}}\ \text{s}$   
 (3)  $2\ \text{s}$                                       (4)  $\frac{\sqrt{3}}{2}\ \text{s}$