

# JEE–MAIN EXAMINATION – JANUARY 2025

(HELD ON WEDNESDAY 29<sup>th</sup> JANUARY 2025)

TIME : 9 : 00 AM TO 12 : 00 NOON

## PHYSICS

### SECTION-A

26. Given below are two statements : one is labelled as **Assertion (A)** and the other is labelled as **Reason (R)**.

**Assertion (A)** : Choke coil is simply a coil having a large inductance but a small resistance. Choke coils are used with fluorescent mercury-tube fittings. If household electric power is directly connected to a mercury tube, the tube will be damaged.

**Reason (R)** : By using the choke coil, the voltage across the tube is reduced by a factor  $\left( R / \sqrt{R^2 + \omega^2 L^2} \right)$ , where  $\omega$  is frequency of the supply across resistor  $R$  and inductor  $L$ . If the choke coil were not used, the voltage across the resistor would be the same as the applied voltage. In the light of the above statements, choose the **most appropriate answer** from the options given below:

- (1) Both (A) and (R) are true but (R) is **not** the correct explanation of (A).
- (2) (A) is false but (R) is true.
- (3) Both (A) and (R) are true and (R) is the correct explanation of (A).
- (4) (A) is true but (R) is false.

27. Two projectiles are fired with same initial speed from same point on ground at angles of  $(45^\circ - \alpha)$  and  $(45^\circ + \alpha)$ , respectively, with the horizontal direction. The ratio of their maximum heights attained is :

- (1)  $\frac{1 - \tan \alpha}{1 + \tan \alpha}$
- (2)  $\frac{1 + \sin \alpha}{1 - \sin \alpha}$
- (3)  $\frac{1 - \sin 2\alpha}{1 + \sin 2\alpha}$
- (4)  $\frac{1 + \sin 2\alpha}{1 - \sin 2\alpha}$

28. An electric dipole of mass  $m$ , charge  $q$ , and length  $l$  is placed in a uniform electric field  $\vec{E} = E_0 \hat{i}$ . When the dipole is rotated slightly from its equilibrium position and released, the time period of its oscillations will be :

- (1)  $\frac{1}{2\pi} \sqrt{\frac{2ml}{qE_0}}$
- (2)  $2\pi \sqrt{\frac{ml}{qE_0}}$
- (3)  $\frac{1}{2\pi} \sqrt{\frac{ml}{2qE_0}}$
- (4)  $2\pi \sqrt{\frac{ml}{2qE_0}}$

29. The pair of physical quantities not having same dimensions is :

- (1) Torque and energy
- (2) Surface tension and impulse
- (3) Angular momentum and Planck's constant
- (4) Pressure and Young's modulus

30. Given below are two statements : one is labelled as **Assertion (A)** and the other is labelled as **Reason (R)**.

**Assertion (A)** : Time period of a simple pendulum is longer at the top of a mountain than that at the base of the mountain.

**Reason (R)** : Time period of a simple pendulum decreases with increasing value of acceleration due to gravity and vice-versa.

In the light of the above statements, choose the **most appropriate answer** from the options given below:

- (1) Both (A) and (R) are true but (R) is **not** the correct explanation of (A).
- (2) Both (A) and (R) are true and (R) is the correct explanation of (A).
- (3) (A) is true but (R) is false.
- (4) (A) is false but (R) is true.

31. The expression given below shows the variation of velocity ( $v$ ) with time ( $t$ ),  $v = At^2 + \frac{Bt}{C+t}$ . The dimension of ABC is :

- (1)  $[M^0L^2T^{-3}]$  (2)  $[M^0L^1T^{-3}]$   
 (3)  $[M^0L^1T^{-2}]$  (4)  $[M^0L^2T^{-2}]$

32. Consider  $I_1$  and  $I_2$  are the currents flowing simultaneously in two nearby coils 1 & 2, respectively. If  $L_1$  = self inductance of coil 1,  $M_{12}$  = mutual inductance of coil 1 with respect to coil 2, then the value of induced emf in coil 1 will be

- (1)  $\varepsilon_1 = -L_1 \frac{dI_1}{dt} + M_{12} \frac{dI_2}{dt}$   
 (2)  $\varepsilon_1 = -L_1 \frac{dI_1}{dt} - M_{12} \frac{dI_1}{dt}$   
 (3)  $\varepsilon_1 = -L_1 \frac{dI_1}{dt} - M_{12} \frac{dI_2}{dt}$   
 (4)  $\varepsilon_1 = -L_1 \frac{dI_2}{dt} - M_{12} \frac{dI_1}{dt}$


33. At the interface between two materials having refractive indices  $n_1$  and  $n_2$ , the critical angle for reflection of an em wave is  $\theta_{1c}$ . The  $n_2$  material is replaced by another material having refractive index  $n_3$ , such that the critical angle at the interface between  $n_1$  and  $n_3$  materials is  $\theta_{2c}$ . If  $n_3 > n_2 > n_1$ ,  $\frac{n_2}{n_3} = \frac{2}{5}$  and  $\sin \theta_{2c} - \sin \theta_{1c} = \frac{1}{2}$ , then  $\theta_{1c}$  is

- (1)  $\sin^{-1}\left(\frac{1}{6n_1}\right)$  (2)  $\sin^{-1}\left(\frac{2}{3n_1}\right)$   
 (3)  $\sin^{-1}\left(\frac{5}{6n_1}\right)$  (4)  $\sin^{-1}\left(\frac{1}{3n_1}\right)$

34. Consider a long straight wire of a circular cross-section (radius  $a$ ) carrying a steady current  $I$ . The current is uniformly distributed across this cross-section. The distances from the centre of the wire's cross-section at which the magnetic field [inside the wire, outside the wire] is half of the maximum possible magnetic field, any where due to the wire, will be

- (1)  $[a/4, 3a/2]$  (2)  $[a/2, 2a]$   
 (3)  $[a/2, 3a]$  (4)  $[a/4, 2a]$

35. As shown below, bob A of a pendulum having massless string of length 'R' is released from  $60^\circ$  to the vertical. It hits another bob B of half the mass that is at rest on a friction less table in the centre. Assuming elastic collision, the magnitude of the velocity of bob A after the collision will be (take  $g$  as acceleration due to gravity)



- (1)  $\frac{1}{3}\sqrt{Rg}$  (2)  $\sqrt{Rg}$   
 (3)  $\frac{4}{3}\sqrt{Rg}$  (4)  $\frac{2}{3}\sqrt{Rg}$

36. Given below are two statements : one is labelled as **Assertion (A)** and the other is labelled as **Reason (R)**.

**Assertion (A)** : Emission of electrons in photoelectric effect can be suppressed by applying a sufficiently negative electron potential to the photoemissive substance.

**Reason (R)** : A negative electric potential, which stops the emission of electrons from the surface of a photoemissive substance, varies linearly with frequency of incident radiation.

In the light of the above statements, choose the **most appropriate answer** from the options given below:

- (1) (A) is false but (R) is true.
- (2) (A) is true but (R) is false.
- (3) Both (A) and (R) are true and (R) is the correct explanation of (A).
- (4) Both (A) and (R) are true but (R) is **not** the correct explanation of (A).


37. A coil of area  $A$  and  $N$  turns is rotating with angular velocity  $\omega$  in a uniform magnetic field  $\vec{B}$  about an axis perpendicular to  $\vec{B}$ . Magnetic flux  $\phi$  and induced emf  $\varepsilon$  across it, at an instant when  $\vec{B}$  is parallel to the plane of coil, are :

- (1)  $\phi = AB, \varepsilon = 0$
- (2)  $\phi = 0, \varepsilon = NAB\omega$
- (3)  $\phi = 0, \varepsilon = 0$
- (4)  $\phi = AB, \varepsilon = NAB\omega$


38. The fractional compression  $\left(\frac{\Delta V}{V}\right)$  of water at the depth of 2.5 km below the sea level is \_\_\_\_\_.  
Given, the Bulk modulus of water  $= 2 \times 10^9 \text{ Nm}^{-2}$ , density of water  $= 10^3 \text{ kg m}^{-3}$ , acceleration due to gravity  $= g = 10 \text{ ms}^{-2}$ .

- (1) 1.75
- (2) 1.0
- (3) 1.5
- (4) 1.25

39. If  $\lambda$  and  $K$  are de Broglie Wavelength and kinetic energy, respectively, of a particle with constant mass. The correct graphical representation for the particle will be :-



40.




For the circuit shown above, equivalent GATE is :

- (1) OR gate
- (2) NOT gate
- (3) AND gate
- (4) NAND gate

41. A body of mass 'm' connected to a massless and unstretchable string goes in verticle circle of radius 'R' under gravity  $g$ . The other end of the string is fixed at the center of circle. If velocity at top of circular path is  $n\sqrt{gR}$ , where,  $n \geq 1$ , then ratio of kinetic energy of the body at bottom to that at top of the circle is

- (1)  $\frac{n}{n+4}$
- (2)  $\frac{n+4}{n}$
- (3)  $\frac{n^2}{n^2+4}$
- (4)  $\frac{n^2+4}{n^2}$

42. Let  $u$  and  $v$  be the distances of the object and the image from a lens of focal length  $f$ . The correct graphical representation of  $u$  and  $v$  for a convex lens when  $|u| > f$ , is



43. Match **List-I** with **List-II**.

	List-I		List-II
(A)	Electric field inside (distance $r > 0$ from center) of a uniformly charged spherical shell with surface charge density $\sigma$ , and radius $R$ .	(I)	$\sigma / \epsilon_0$
(B)	Electric field at distance $r > 0$ from a uniformly charged infinite plane sheet with surface charge density $\sigma$ .	(II)	$\sigma / 2\epsilon_0$
(C)	Electric field outside (distance $r > 0$ from center) of a uniformly charged spherical shell with surface charge density $\sigma$ , and radius $R$ .	(III)	0
(D)	Electric field between 2 oppositely charged infinite plane parallel sheets with uniform surface charge density $\sigma$ .	(IV)	$\frac{\sigma}{\epsilon_0 r^2}$

Choose the **correct** answer from the options given below :

- (1) (A)-(IV), (B)-(I), (C)-(III), (D)-(II)
- (2) (A)-(IV), (B)-(II), (C)-(III), (D)-(I)
- (3) (A)-(II), (B)-(I), (C)-(IV), (D)-(III)
- (4) (A)-(III), (B)-(II), (C)-(IV), (D)-(I)

44. The workdone in an adiabatic change in an ideal gas depends upon only :

- (1) change in its pressure
- (2) change in its specific heat
- (3) change in its volume
- (4) change in its temperature

45. Given below are two statements : one is labelled as **Assertion (A)** and other is labelled as **Reason (R)**.

**Assertion (A)** : Electromagnetic waves carry energy but not momentum.

**Reason (R)** : Mass of a photon is zero.

In the light of the above statements, choose the **most appropriate answer** from the options given below :

- (1) (A) is true but (R) is false.
- (2) (A) is false but (R) is true.
- (3) Both (A) and (R) are true but (R) is **not** the correct explanation of (A).
- (4) Both (A) and (R) are true and (R) is the correct explanation of (A).

**SECTION-B**


46. The coordinates of a particle with respect to origin in a given reference frame is (1, 1, 1) meters. If a force of  $\vec{F} = \hat{i} - \hat{j} + \hat{k}$  acts on the particle, then the magnitude of torque (with respect to origin) in z-direction is \_\_\_\_\_.

47. A container of fixed volume contains a gas at  $27^\circ\text{C}$ . To double the pressure of the gas, the temperature of gas should be raised to \_\_\_\_\_  $^\circ\text{C}$ .

48. Two light beams fall on a transparent material block at point 1 and 2 with angle  $\theta_1$  and  $\theta_2$ , respectively, as shown in figure. After refraction, the beams intersect at point 3 which is exactly on the interface at other end of the block. Given : the distance between 1 and 2,  $d = 4\sqrt{3}$  cm and

$$\theta_1 = \theta_2 = \cos^{-1} \left( \frac{n_2}{2n_1} \right), \text{ where refractive index of}$$

the block  $n_2 >$  refractive index of the outside medium  $n_1$ , then the thickness of the block is \_\_\_\_\_ cm.



49. In a hydraulic lift, the surface area of the input piston is  $6 \text{ cm}^2$  and that of the output piston is  $1500 \text{ cm}^2$ . If  $100 \text{ N}$  force is applied to the input piston to raise the output piston by  $20 \text{ cm}$ , then the work done is \_\_\_\_\_ kJ.
50. The maximum speed of a boat in still water is  $27 \text{ km/h}$ . Now this boat is moving downstream in a river flowing at  $9 \text{ km/h}$ . A man in the boat throws a ball vertically upwards with speed of  $10 \text{ m/s}$ . Range of the ball as observed by an observer at rest on the river bank, is \_\_\_\_\_ cm.  
(Take  $g = 10 \text{ m/s}^2$ )